



**NATIONAL SENIOR
CERTIFICATE/
NASIONALE SENIOR
SERTIFIKAAT**

GRADE/GRAAD 12

JUNE/JUNIE 2022

**PHYSICAL SCIENCES: CHEMISTRY P2
MARKING GUIDELINE/
FISIESE WETENSKAPPE: CHEMIE V2
NASIENRIGLYN**

MARKS/PUNTE: 150

This marking guideline consists of 13 pages./
Hierdie nasienriglyn bestaan uit 13 bladsye.

QUESTION/VRAAG 1

- 1.1 A ✓✓ (2)
- 1.2 B ✓✓ (2)
- 1.3 A ✓✓ (2)
- 1.4 C ✓✓ (2)
- 1.5 B ✓✓ (2)
- 1.6 D ✓✓ (2)
- 1.7 B ✓✓ (2)
- 1.8 A ✓✓ (2)
- 1.9 B ✓✓ (2)
- 1.10 D ✓✓ (2)
- [20]**

QUESTION/VRAAG 2

2.1 2.1.1 Alkyne ✓ (1)

2.1.2 Haloalkane ✓ (1)

2.2 2.2.1 E ✓ (1)


2.2.2 D ✓ (1)

2.3 2.3.1 Compound that contains carbon and hydrogen atoms only. ✓✓ (2 or 0)
Verbinding wat slegs koolstof- en waterstof-atome bevat. (2 of 0) (2)

2.3.2 UNSATURATED ✓ Contains triple bond ✓/multiple bonds (between the C-atoms in die hydrocarbon chain)
ONVERSADIG Bevat 'n drievoudige binding/ meervoudige bindings (tussen C-atome in die koolwaterstofketting) (2)

2.3.3 6-ethyl ✓-2-methyl ✓ oct-4-yne ✓ / 6-ethyl-2-methyl-4-octyne
6-etiel-2-metielokt-4-yne / 6-etiel-2-metiel-4-oktyn (3)

2.4




Marking criteria/Nasienkriteria:

- Whole structure correct/*Hele struktuur korrek: (2/2)*
- Only functional group correct
Slegs funksionele groep korrek Max./Maks. (1/2)

(2)

2.5 2.5.1 Butanal ✓✓ (2)

2.5.2



Marking criteria/Nasienriglyne

- Whole structure correct/*Hele struktuur korrek: (2/2)*
- Only functional group correct
Slegs funksionele groep korrek Max./Maks. (1/2)

(2)

2.6 2.6.1 Combustion ✓/Oxidation
Verbranding / Oksidasie (1)

2.6.2 $2 \text{C}_6\text{H}_{14} + 19 \text{O}_2 \checkmark \rightarrow 12 \text{CO}_2 + 14 \text{H}_2\text{O} \checkmark$ (✓ Balancing/ *balansering*) (3)

2.6.3 Compound A reacts exothermically with oxygen / releases heat when it reacts with oxygen. ✓
Verbinding A reageer eksotermies met suurstof / hitte word vrygestel wanneer dit met suurstof reageer. (1)

[22]

QUESTION/VRAAG 3

- 3.1 3.1.1 Boiling point is the temperature ✓ at which the vapour pressure of a liquid equals the atmospheric pressure. ✓
Kookpunt is die temperatuur waarby die dampdruk van 'n vloeistof gelyk is aan die atmosferiese druk. (2)
- 3.1.2 Functional group/ ✓ Homologous series/ Type of intermolecular forces.
Funksionele groep / Homoloë reeks / Tipe intermolekulêre kragte. (1)
- 3.1.3 London forces ✓ / induced dipole forces / dispersion forces.
Londonkragte / geïnduseerde dipool kragte / verspreidingskragte. (1)
- 3.1.4 **C** has hydrogen bonds ✓ (in addition to London forces)
B has dipole-dipole forces ✓ (in addition to London forces)
 Hydrogen bonds are stronger than dipole-dipole forces ✓
 More energy is needed to overcome intermolecular forces in **C** ✓
***C** het waterstofbinding (bykomend tot Londonkragte)*
***B** het dipool-dipoolkragte (bykomend tot Londonkragte)*
Waterstofbindings is sterker as die dipool-dipoolkragte
*Meer energie word benodig om die intermolekulêre kragte te oorkom in **C***
- OR/OF**
- C** has hydrogen bonds ✓ (in addition to London forces)
B has dipole-dipole forces ✓ (in addition to London forces)
 Dipole-dipole forces are weaker than hydrogen bonds ✓
 Less energy is needed to overcome intermolecular forces in **B** ✓
***C** het waterstofbinding (bykomend tot Londonkragte)*
***B** het dipool-dipoolkragte (bykomend tot Londonkragte)*
Dipool-dipool is swakker as die waterstofbindings
*Minder energie word benodig om die intermolekulêre kragte te oorkom in **B*** (4)
- 3.2 3.2.1 Vapour pressure is the pressure exerted by a vapour ✓ in equilibrium with its liquid in a closed container. ✓
Dampdruk is die druk uitgeoefen deur 'n damp in ewewig met sy vloeistof in 'n geslote sisteem. (2)
- 3.2.2 $p_1 = 100$ (kPa) ✓ / 101,3 (kPa) / 1 atmosphere / *atmosfeer* (1)
- 3.2.3 Gas ✓ **A** is above its boiling point / *Bo **A** se kookpunt* ✓ (2)
- 3.2.4 LOWER THAN/LAER AS ✓ (1)

- 3.2.5 Compound C only reached its boiling point at 117,7 °C where its vapour pressure will equal 101,3 kPa. ✓✓
Verbinding C bereik eers sy kookpunt by 117,7 °C waar sy dampdruk eers gelyk aan 101,3 kPa gaan wees.


(2)
[16]

QUESTION/VRAAG 4

- 4.1 Substitution ✓/Hydrolysis (of haloalkanes)
Substitusie / Hidrolise (van haloalkane) (1)
- 4.2 $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ ✓✓ (2)
- 4.3 Primary alcohol ✓
The carbon atom that contains the hydroxyl group (-OH) is bonded to one other carbon atom only. ✓
Primêre alkohol
Die koolstof-atoom wat die hidroksielgroep (-OH) bevat is verbind aan slegs een ander koolstof-atoom. (2)
- 4.4 Esterification/Condensation ✓
Esterifikasie / Kondensasie (1)
- 4.5 Butyl ethanoate / *Butiel-etanoaat* ✓✓



Marking criteria/Nasienkriteria:

- Functional group/*Funksionele groep.* ✓ (1/2)
- Whole structure correct/
Hele struktuur korrek ✓ (2/2)



- 4.6 Dehydration / *Dehidrasie* ✓ (1)
- 4.7 (concentrated / *gekonsentreerde*) H_2SO_4 ✓ (1)
- 4.8 Compounds with same molecular formula ✓ but different positions of the functional group ✓/side chains/substituent on the parent chain./
Verbindings met dieselfde molekulêre formule maar verskillende posisies van die funksionele groep/sykettings/substituent op die stamketting. (2)

4.9

**Marking criteria/ Nasienkriteria****Reactants / Reaktanse**

- Organic molecule correct/ *Organiese molekule korrek* ✓
- KOH/NaOH ✓

Products / Produkte

- **Organic molecule / Organiese molekule**
- Functional group/*Funksionele groep.* ✓ (1/2)
- Whole structure correct/
Hele struktuur korrek ✓ (2/2)

Inorganic products / Anorganiese produkte

- KBr/NaBr ✓
- H₂O ✓

(6)
[20]

QUESTION 5/VRAAG 5

- 5.1 Change in concentration ✓ per unit time. ✓ / Amount of product formed/reactant used up per unit time.
Verandering in konsentrasie per eenheid tyd / Hoeveelheid produk gevorm / reaktanse opgebruik per eenheidstyd.

OR/OF

Rate of change in concentration **(2 or 0)**
Tempo van verandering in konsentrasie (2 of 0) (2)

- 5.2 Concentration ✓ and a catalyst. ✓ /
Konsentrasie en 'n katalisator. (2)

- 5.3 5.3.1 Concentration (of H₂O₂) decreases / *Konsentrasie (van H₂O₂) verlaag* ✓✓ (2)

- 5.3.2 $t = 5\,000\text{(s)}$ ✓ (1)

- 5.3.3 Rate/tempo = $-\Delta c/\Delta t = - (1,0 - 0) / (0 - 5\,000)$ ✓
 $= 2 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3}\cdot\text{s}^{-1}$ ✓ (3)

- 5.3.4 $\Delta c = 0,8 \text{ mol}\cdot\text{dm}^{-3}$

$$\begin{aligned} n(\text{H}_2\text{O}_2) &= cV \\ &= 0,8 \times (0,15) \text{ ✓} \\ &= 0,12 \text{ mol} \end{aligned}$$

$$n(\text{H}_2\text{O}_2) = n(\text{O}_2) = 0,12 \text{ mol} \text{ ✓ (Ratio / verhouding)}$$

$$n = V/V_m \text{ ✓}$$

$$0,12 = V/25\,000 \text{ ✓}$$

$$V = 3\,000 \text{ cm}^3 \text{ ✓} / (3 \text{ dm}^3) \quad (5)$$

- 5.4 5.4.1 Decrease / *Verlaag* ✓ (1)

- 5.4.2 Remains the same / *Bly dieselfde* ✓ (1)

- 5.5 5.5.1 Minimum energy required for a reaction to take place. ✓✓ /
Minimum energie benodig vir 'n reaksie om plaas te vind. (2)

- 5.5.2 Particles with sufficient kinetic energy to react. ✓ /
Deeltjies met genoeg kinetiese energie om te reageer. (1)

5.5.3 E₂ ✓

Catalyst lowers the activation energy ✓

More particles have sufficient kinetic energy to react ✓

More effective collisions per unit time ✓ / Frequency of effective collisions increases

'n Katalisator verlaag die aktiveringsenergie.

Meer deeltjies het genoeg kinetiese energie om te reageer

Meer effektiewe botsings per eenheid tyd / Frekwensies van die effektiewe botsings neem toe

(4)
[24]

QUESTION 6/VRAAG 6

6.1 6.1.1 Reversible ✓ (reaction) / Omkeerbare (reaksie) (1)

6.1.2 HIGHER THAN / HOËR AS ✓✓ (2)

6.1.3 EQUAL TO / GELYK AAN ✓ (1)

6.2 **OPTION 1/ OPSIE 1 : MOLE OPTION / MOL OPSIE**

Marking Criteria/ Nasienkriteria:

- Divide by 17 to calculate $n(\text{NH}_3)_{\text{equilibrium}}$. ✓
Deel deur 17 om $n(\text{NH}_3)_{\text{ewewig}}$ te bereken.
- $\Delta n \text{ NH}_3$ ✓
- Use mole ratio $\text{N}_2:\text{H}_2:\text{NH}_3$ / *Gebruik mol verhouding $\text{N}_2:\text{H}_2:\text{NH}_3$* ✓
- $n_{\text{equilibrium}}$ / *ewewig N_2 and/en H_2* ✓
- Divide 2 dm^3 all $n_{\text{equilibrium}}$ / *Deel deur 2 dm^3 in alle n_{ewewig}* ✓
- Correct K_c expression / *Korrekte K_c uitdrukking.* ✓
- Substitution into K_c expression ✓ / *Vervanging in K_c uitdrukking.*
- Final answer / *Finale antwoord (0,41)* ✓

$$n(\text{NH}_3) = m/M = 41,48/17 \checkmark = 2,44 \text{ mol}$$

	N_2	3 H_2	2 NH_3	
n_i	4,88	6,18	0	
Δn	1,22	3,66	2,44 ✓	Ratio / verhouding
n_e	3,66	2,52 ✓	2,44 ✓	
C_e	1,83	1,26	1,22 ✓	(Div/by deel met 2 dm^3)

$$\begin{aligned} K_c &= \frac{[\text{NH}_3]^2}{[\text{N}_2] \cdot [\text{H}_2]^3} \checkmark \\ &= \frac{1,22^2}{1,83 \times 1,26^3} \checkmark \\ &= 0,41 \checkmark \end{aligned}$$

OPTION 2: CONCENTRATION / OPSIE 2: KONSENTRASIE**Marking criteria/ Nasienkriteria:**

- Calculate $c(\text{NH}_3)_{\text{equilibrium}}$. ✓
Bereken $c(\text{NH}_3)_{\text{ewewig}}$
- $\Delta c \text{ NH}_3$ ✓
- $c_i \text{ N}_2$ and/en H_2 ✓
- Use conc. ratio $\text{N}_2:\text{H}_2:\text{NH}_3$ /Gebruik gekonsen. verhouding $\text{N}_2:\text{H}_2:\text{NH}_3$ ✓
- c equilibrium /ewewig N_2 and/en H_2 ✓
- Correct K_c expression/Korrekte K_c uitdrukking. ✓
- Substitution into K_c expression ✓/Vervanging in K_c uitdrukking.
- Final answer/Finale antwoord (0,41) ✓

$$n(\text{NH}_3) = m/M = 41,48/17 = 2,44 \text{ mol}$$

$$c_e(\text{NH}_3) = n/V = 2,44/2 \quad \checkmark \quad = 1,22 \text{ mol}\cdot\text{dm}^{-3}$$

$$c_i(\text{N}_2) = n/V = 4,88/2 = 2,44 \text{ mol}\cdot\text{dm}^{-3}$$

$$c_i(\text{H}_2) = n/V = 6,18/2 = 3,09 \text{ mol}\cdot\text{dm}^{-3}$$

	N_2	3 H_2	2 NH_3	
c_i	2,44	3,09	0	
Δc	0,61	1,83	1,22 ✓	Ratio / verhouding
c_e	<u>1,83</u>	<u>1,26</u> ✓	1,22 ✓	

$$K_c = [\text{NH}_3]^2/[\text{N}_2]\cdot[\text{H}_2]^3 \quad \checkmark$$

$$= 1,22^2/1,83 \times 1,26^3 \quad \checkmark$$

$$= 0,41 \quad \checkmark$$

(8)

6.3 Temperature / *Temperatuur* ✓

(1)

6.4 **A** ✓At a given temperature the yield of NH_3 is the highest ✓ (in graph **A**)

Increase pressure favours reaction which produces less gas moles ✓

Forward reaction is favoured ✓

*By 'n gegewe temperatuur is die opbrengs van NH_3 die hoogste (in grafiek **A**)**Toename in druk bevoordeel die reaksie wat die minste gas mol produseer**Voorwaartse reaksie word bevoordeel*

(4)

6.5 6.5.1 Increase / *Toeneem* ✓

(1)

6.5.2 No effect / *Geen effek* ✓

(1)

6.5.3 Decrease/ *Afneem* ✓

(1)

[20]

QUESTION 7/VRAAG 7

7.1 7.1.1 An acid is a proton (H^+ -ion) donor ✓✓
'n Suur is 'n proton (H^+ -ioon) -skenker (2)

7.1.2 H_2O ✓ and / en $H_2SO_4^-$ ✓ (2)

7.1.3 H_2O or/of HSO_4^- (Any ONE / Enige EEN ✓) (1)

7.2 7.2.1 WEAK ACID ✓ Low K_a value/ $K_a < 1$ ✓
SWAKSUUR Lae K_a -waarde / $K_a < 1$ (2)

7.2.2 H_2SO_4 ✓ (1)

7.2.3 $pH = -\log [H_3O^+]$ ✓

$$3 = -\log [H_3O^+] \quad \checkmark$$

$$[H_3O^+] = 1 \times 10^{-3} \text{ mol}\cdot\text{dm}^{-3}$$

$$[H_2SO_4] = \frac{1}{2} \times 10^{-3} \quad \checkmark$$

$$= 5 \times 10^{-4} \text{ mol}\cdot\text{dm}^{-3} \quad \checkmark \quad (4)$$

7.2.4 $CO_3^{2-} + H_2O \quad \checkmark \quad \rightarrow \quad HCO_3^- + OH^- \quad \checkmark$



(Excess) OH^- formed / (oormaat) OH^- word geproduseer ✓ (3)

7.3.1 It is a solution of known concentration ✓✓
Dit is 'n oplossing van bekende konsentrasie (2)

7.3.2	<p><u>OPTION 1 / OPSIE 1:</u> $n = m/M$ $n = 1,74/58 \quad \checkmark$ $= 0,03 \text{ mol}$ $c = n/V$ $= 0,03 / 0,2 \quad \checkmark$ $= 0,15 \text{ mol}\cdot\text{dm}^{-3}$</p>	<p><u>OPTION 2 / OPSIE 2:</u> $c = m/MV$ $c = 1,74 / (58)(0,2) \quad \checkmark \checkmark$ $= 0,15 \text{ mol}\cdot\text{dm}^{-3}$</p>
	<p><u>OPTION 3 / OPSIE 3:</u> $n = cV$ $= (0,15)(0,2) \quad \checkmark$ $n = 0,3 \text{ mol}$ $m = nM$ $= (0,3)(58) \quad \checkmark$ $= 1,74 \text{ g}$</p>	<p><u>OPTION 4 / OPSIE 4:</u> $m = cMV$ $= (0,15)(58)(0,2) \quad \checkmark \checkmark$ $m = 1,74 \text{ g}$</p>

(2)

7.3.3 **Marking guideline / Nasienriglyn**

- Calculating mole for $\text{Mg}(\text{OH})_2$ / Bereken mol vir $\text{Mg}(\text{OH})_2$
- Calculating the diluted concentration for HCl / Bereken die verdunde konsentrasie van HCl
- Calculating the total mole for HCl / Bereken die totale mol van HCl
- Calculating the reacted mole for HCl / Bereken die mol van HCl wat gereageer het
- Determining the remaining mole for HCl / Bepaal die mol van HCl wat oorbly
- Use of formula $c = n/V$ in calculating the concentration of excess ions / Gebruik formule $c = n/V$ om die konsentrasie van die oormaat ione te bereken
- Substituting into / Vervanging in $c = n/V$
- Final answer/ Finale antwoord

$$\begin{aligned} n[\text{Mg}(\text{OH})_2] &= cV \\ &= 0,15 \times 0,04 \checkmark \\ &= 0,006 \text{ mol} \end{aligned}$$

$$\begin{aligned} c_1V_1 &= c_2V_2 \\ 5 \times 10 &= c_2(100) \checkmark \\ c_2 &= 0,5 \text{ mol}\cdot\text{dm}^{-3} \end{aligned} \quad \text{OR/OF}$$

$$\begin{aligned} n &= cV \\ &= (10)5 \times 10^{-3} \\ n &= 0,05 \text{ mol} \\ c &= \frac{n}{V} \\ &= \frac{0,05}{0,1} \\ c &= 0,5 \text{ mol}\cdot\text{dm}^{-3} \end{aligned} \quad \checkmark$$

$$\begin{aligned} n(\text{HCl}) &= cV \\ &= 0,5 \times 0,05 \checkmark \\ &= 0,025 \text{ mol} \end{aligned}$$

$$n(\text{HCl}) \text{ reacting} = 2 \times 0,006 \checkmark = 0,012 \text{ mol}$$

$$n(\text{HCl}) \text{ remaining} = 0,025 - 0,012 \checkmark\checkmark = 0,013 \text{ mol}$$

$$c(\text{HCl}) = n/V \checkmark = 0,013 / (0,09) \checkmark = 0,14 \text{ mol}\cdot\text{dm}^{-3} \checkmark$$

(9)
[28]**TOTAL/TOTAAL: 150**